

Emergence of triplet orbital pairing and non-Abelian states in ultracold multi-orbital optical lattices with quadratic band touching

Jian-Hua Jiang^{1,*}

¹*Department of Condensed Matter Physics, Weizmann Institute of Science, Rehovot 76100, Israel*
(Dated: December 6, 2011)

It is found that all the *singlet orbital pairing* instabilities are *absent* in a class of spin-polarized multi-orbital systems with quadratic band touching, which opens the way for *triplet orbital pairing* order. The ground states are found to be *non-Abelian* states with *p*-wave orbital pairing in checkerboard (away from 1/2 filling) and kagome (above 1/3 filling) lattices with *isotropic* attractive interaction which can be realized in ultracold multi-orbital optical lattices. The special property of such systems is generalized to more classes of multi-orbital systems, where the fully-gapped *non-Abelian* states are possibly the ground states. Those findings are helpful in achieving topological quantum computation.

PACS numbers: 67.85.Lm, 03.67.Lx, 71.10.Pm, 03.65.Vf

Introduction.— Excitations obeying non-Abelian statistics can emerge in interacting many-fermion systems[1, 2]. One known prototype is the fully-gapped *p*-wave superconducting (SC)/superfluid (SF) state [3]. More generally, any fully-gapped SC/SF state with *odd* Chern number is a *non-Abelian* state, as there is one topologically protected zero-energy Majorana bound state in each quantized vortex and the braiding of the Majorana fermions leads to the non-Abelian statistics[4]. The search for systems with non-Abelian excitations is desirable not only in the sense that they can be used to realize topological quantum computation (TQC) [2, 5–7], but also that they have nontrivial ground states which are characterized by topological orders[8].

Except in a few cases[9–13], triplet pairing is crucial to the emergence of non-Abelian states. However, in reality triplet pairing is scarce, whereas singlet pairing prevails. One of the reason is that the interaction between fermions with opposite spin (e.g., on-site interaction) is stronger than that between electrons with the same spin (e.g., nearest neighbor interaction) due to Pauli exclusion. In spinless (spin-polarized) fermionic systems, the situation is different. In lattice systems with a single orbit (site) in an unit cell, only the triplet pairing is possible. However, in multi-orbital systems, where pseudo-spin denotes the orbital degree of freedom, the singlet (inter-orbital) pairing prevails due to similar reasons.

In this Letter, we propose a scenario that suppresses *all* the *singlet* (inter-orbital) pairing instabilities and opens the way to the *triplet* orbital pairing, regardless of the relative strength of the pairing interaction of the two. The concerned systems have a single quadratic band touching (QBT) protected by time-reversal and space-inversion symmetry[14], while the Fermi level is above or below the QBT point. The unique property of such systems is that the pseudo-spin polarization on the Fermi surface has a winding number of ± 2 . Due to such winding as well as the time-reversal symmetry, the $\mathbf{k}$ and $-\mathbf{k}$ states on the Fermi surface has the same pseudo-spin

polarization (pseudo-spin polarization winds one period when $\mathbf{k}$ winds to $-\mathbf{k}$), while the states with opposite pseudo-spin is well below the Fermi surface. Hence there is *no* singlet orbital pairing instability in the weak pairing regime, which opens a way to *triplet orbital pairing orders* in multi-orbital systems. For concreteness, we study two systems with a single QBT: the checkerboard (away from 1/2 filling) and kagome (above 1/3 filling) lattices with *isotropic* attractive interaction. It is found that the ground states in those systems are *non-Abelian* states with *p*-wave orbital pairing which is promising for TQC. Furthermore, the special property of such systems is generalized to more classes of multi-orbital systems, where the *non-Abelian* states which have fully-gapped Fermi surface to gain more condensation energy are possibly the ground states[15, 16].

Ultracold fermionic atom/polar-molecule gases with tunable interaction through Feshbach resonance[17, 18] and other techniques[19] offers a lot of advantages in realizing generic interacting many-fermion systems[20]. Especially, in polar-molecule systems the combination of microwave excitation with dipole-dipole interactions enables a variety of effective inter-molecule interactions in a designable fashion and with significant strength to achieve observable emergent phases[19]. The good controllability and emerging new detection techniques also enable them to be an ideal platform to achieve TQC[7], given that the non-Abelian states can be realized. We show that the systems with non-Abelian ground states proposed in this Letter can be realized in spin-polarized ultracold fermionic systems in multi-orbital optical lattices[14, 21, 25, 26]. With the emerging technology advancements in multi-orbital optical lattices[21] and ultracold fermionic atom/polar-molecule gases[19], the proposed systems are helpful in achieving TQC.

Quadratic band touching in checkerboard and kagome lattices.— The checkerboard lattice is depicted in Fig. 1(a). In each unit cell there are two sites labeled

as red (circle) and blue (square) dots in the figure. Allowing one orbit in each site, for spin-polarized fermions, the Hamiltonian takes the form

$$H = - \sum_{\langle i\sigma, j\sigma' \rangle} t_{i\sigma, j\sigma'} (c_{i\sigma}^\dagger c_{j\sigma'} + H.c.) + H_{\text{int}}. \quad (1)$$

Here i and j are the indices of the unit cells, while pseudo-spins $(\sigma, \sigma' = \uparrow, \downarrow)$ denote the two different orbits in each unit cell. $\langle i\sigma, j\sigma' \rangle$ restricts the summation to the nearest and next nearest neighbors. The system is engineered in such a way that the hopping amplitude in x and y directions between red (blue) sites are t' [solid links in Fig. 1(a)] and t'' [dotted links] (t'' and t') respectively. The hopping between red and blue sites are t [dashed links]. Accordingly, the free Hamiltonian is $H_0 = \sum_{\mathbf{k}} \psi_{\mathbf{k}}^\dagger \mathcal{H}_0(\mathbf{k}) \psi_{\mathbf{k}}$ with $\psi_{\mathbf{k}} = (c_{\mathbf{k}\uparrow}, c_{\mathbf{k}\downarrow})^T$, where

$$\mathcal{H}_0(\mathbf{k}) = h_0(\mathbf{k})\sigma_0 + \mathbf{h}(\mathbf{k}) \cdot \hat{\sigma} \quad (2)$$

where $\hat{\sigma}_0$ denotes the 2×2 identity matrix and $\hat{\sigma}$ is the Pauli matrix vector. $h_0(\mathbf{k}) = -2t_0(\cos \bar{k}_x + \cos \bar{k}_y)$, $h_z(\mathbf{k}) = -2t_z(\cos \bar{k}_x - \cos \bar{k}_y)$, and $h_x(\mathbf{k}) = 8t_x \cos \frac{\bar{k}_x}{2} \cos \frac{\bar{k}_y}{2}$ with $t_0 = (t' + t'')/2$, $t_z = (t' - t'')/2$, $t_x = -t/2$. $h_y(\mathbf{k}) = 0$ due to the time-reversal and space-inversion symmetry[22]. Due to the symmetry, the single QBT point can only be at a time-reversal invariant momentum $\mathbf{K} = -\mathbf{K}$. The QBT is a $\mathbf{k}$ -space vortex which is topologically stable[14]. In checkerboard lattice the two bands touch quadratically [Fig. 1(c)] at $\mathbf{K} = (\pi, \pi)$ [22]. In the vicinity of $\mathbf{K}$, one has $h_0(\mathbf{k}) = t_0 k^2$, $h_z(\mathbf{k}) = t_z(k_x^2 - k_y^2)$, $h_x(\mathbf{k}) = 2t_x k_x k_y$, and $h_y(\mathbf{k}) = 0$, where $\mathbf{k} = \bar{\mathbf{k}} - \mathbf{K}$. As $\mathbf{K}$ is a time-reversal invariant momentum, pairing is between the $\mathbf{k}$ and $-\mathbf{k}$ states. The spectrum is $\varepsilon_{\mathbf{k}\pm} = t_0 k^2 \pm k^2 \sqrt{t_z^2 \cos^2(2\theta_{\mathbf{k}}) + t_x^2 \sin^2(2\theta_{\mathbf{k}})}$ with $\theta_{\mathbf{k}} = \text{Arg}[k_x + ik_y]$. At half-filling, the Fermi level is at the QBT point. Away from it only one band crosses the Fermi level [Fig. 1(c)] when $|t_0| \leq |t_z|, |t_x|$. A characteristic of such systems is that the pseudo-spin polarization on the Fermi surface has a winding number of ± 2 . The winding number is

$$N_w = \frac{1}{2\pi} \oint_{\text{FS}} d\phi_{\mathbf{k}}, \quad (3)$$

where FS stands for the Fermi surface and $\phi_{\mathbf{k}} = \text{Arg}[h_z(\mathbf{k}) + ih_x(\mathbf{k})]$ is the direction of the pseudo-spin polarization in the z - x plane. Winding number $N_w = 2\text{sgn}(t_x t_z) = \pm 2$ as well as the time-reversal symmetry guarantee that the pseudo-spin polarization at $\mathbf{k}$ is the same as that at $-\mathbf{k}$ on the Fermi surface [Fig. 1(d)].

In the kagome lattice, there are three different sites in each unit cell, labeled as red (circle), blue (square) and green (triangle) in Fig. 1(b), which we denote as 1, 2, and 3. With only the nearest neighbor hopping, the free Hamiltonian can be written as $H_0 = \sum_{\mathbf{k}} \psi_{\mathbf{k}}^\dagger \mathcal{H}_{\text{kgm}}(\mathbf{k}) \psi_{\mathbf{k}}$

where $\psi_{\mathbf{k}} = (c_{\mathbf{k}1}, c_{\mathbf{k}2}, c_{\mathbf{k}3})^T$ and

$$\mathcal{H}_{\text{kgm}}(\mathbf{k}) = -2t \begin{pmatrix} 0 & \cos \frac{k_{12}}{2} & \cos \frac{k_{13}}{2} \\ \cos \frac{k_{12}}{2} & 0 & \cos \frac{k_{23}}{2} \\ \cos \frac{k_{13}}{2} & \cos \frac{k_{23}}{2} & 0 \end{pmatrix}$$

with t being the hopping amplitude. $k_{ij} = \mathbf{k} \cdot \mathbf{n}_{ij}$ for $i, j = 1, 2, 3$ with $\mathbf{n}_{12} = (1, 0)$, $\mathbf{n}_{13} = (\frac{1}{2}, \frac{\sqrt{3}}{2})$ and $\mathbf{n}_{23} = \mathbf{n}_{13} - \mathbf{n}_{12}$. The spectrum is $E_{\mathbf{k}0} = 2t$, $E_{\mathbf{k}\pm} = -t \pm t \sqrt{4(\cos^2 \frac{k_{12}}{2} + \cos^2 \frac{k_{13}}{2} + \cos^2 \frac{k_{23}}{2}) - 3}$. In kagome lattice the $+$ and 0 bands touch quadratically at $\mathbf{K} = (0, 0)$. Around $\mathbf{K}$, the spectrum is approximately $E_{\mathbf{k}+} = 2t - \frac{1}{4}tk^2$, $E_{\mathbf{k}0} = 2t$, and $E_{\mathbf{k}-} = -4t + \frac{1}{4}tk^2$. For $t < 0$ and not much above $1/3$ filling (the results are similar for $t > 0$ below $2/3$ filling), only the $+$ band crosses the Fermi surface. Projecting out the $-$ band which is far above, one obtains an effective Hamiltonian in the form of Eq. (2) with $h_0(\mathbf{k}) = \frac{1}{8}|t|k^2$, $h_z(\mathbf{k}) = \frac{1}{8}|t|(k_x^2 - k_y^2)$, $h_x(\mathbf{k}) = \frac{1}{4}|t|k_x k_y$, and $h_y(\mathbf{k}) = 0$, where the pseudo-spin up and down states are defined as $|\uparrow\rangle = \frac{1}{\sqrt{2}}(1, -1, 0)^T$ and $|\downarrow\rangle = \frac{1}{\sqrt{6}}(1, 1, -2)^T$ respectively.




FIG. 1. (Color online) (a) Checkerboard and (b) kagome lattices. In checkerboard lattice, the hopping amplitude along the blue (solid), red (dotted), and green (dashed) links are t' , t'' , and t respectively. (c) Band structure and band filling near half-filling with $t_x = t_z$ in the checkerboard lattice. $\mathbf{k}$ is measured from the QBT point $\mathbf{K}$. (d) Direction of the pseudo-spin field (h_x, h_z) (also represents the pseudo-spin polarization direction) on the Fermi surface of a QBT system.

Nonexistence of singlet pairing instability in QBT systems.— Diagonalizing the Bogoliubov-de Gennes (BdG) Hamiltonian with singlet pairing, $H_{\text{int}}^{\text{sp}} = -\frac{1}{2} \sum_{\mathbf{k}} \Delta_s(\mathbf{k}) (c_{\mathbf{k}\uparrow}^\dagger c_{-\mathbf{k}\downarrow}^\dagger - c_{\mathbf{k}\downarrow}^\dagger c_{-\mathbf{k}\uparrow}^\dagger) + H.c.$, in QBT systems, one finds that the spectrum is *gapless*. The gapless feature is most transparent when $h_0(\mathbf{k}) = 0$, where the spectrum is $\pm [\varepsilon_{\mathbf{k}\pm} - \sqrt{\mu^2 + |\Delta_s(\mathbf{k})|^2}]$ with μ being the chemical potential. For such spectrum, there is *no* pairing instability in the weak pairing regime.

Interaction and p -wave pairing.— Consider the triplet

pairing instabilities due to *isotropic* attractions between fermions at nearest and next nearest neighbor sites,

$$H_{\text{int}} = -\frac{1}{2} \sum_{\langle i\sigma, j\sigma' \rangle} V_{\sigma\sigma'} n_{i\sigma} n_{j\sigma'}. \quad (4)$$

The physical realization of such interaction will be discussed in the end of the paper. In checkerboard lattices, we denote $V_{\uparrow\downarrow} = V$ and $V_{\uparrow\uparrow} = V_{\downarrow\downarrow} = U$. Using a Hubbard-Stratonovich decoupling with BCS pairing and ignoring superconducting fluctuations, one gets

$$H_{\text{int}}^{\text{MF}} = -\frac{1}{2} \sum_{\mathbf{k}\nu} [\Delta_\nu(\mathbf{k}) d_\nu(\mathbf{k}) + H.c.] + \frac{1}{2} E_0, \quad (5)$$

where $d_{x/y}(\mathbf{k}) = c_{\mathbf{k}\uparrow}^\dagger c_{-\mathbf{k}\uparrow}^\dagger \mp c_{\mathbf{k}\downarrow}^\dagger c_{-\mathbf{k}\downarrow}^\dagger$, $d_z(\mathbf{k}) = c_{\mathbf{k}\uparrow}^\dagger c_{-\mathbf{k}\downarrow}^\dagger + c_{\mathbf{k}\downarrow}^\dagger c_{-\mathbf{k}\uparrow}^\dagger$, and $E_0 = \sum_{\mathbf{k}\nu} \Delta_\nu(\mathbf{k}) \langle d_\nu(\mathbf{k}) \rangle$. The triplet pairing is $\Delta_\nu(\mathbf{k}) = V_\nu \sum_{\mathbf{p}} S(\mathbf{k}, \mathbf{p}) \langle d_\nu^\dagger(\mathbf{p}) \rangle$, where $V_x = V_y = U$, $V_z = V/2$, and $S(\mathbf{k}, \mathbf{p}) \simeq \frac{1}{2}(k_+ p_- + k_- p_+)$ with $k_\pm = k_x \pm i k_y$. Hence the pairing is *p*-wave type.

For kagome lattice with *isotropic* nearest neighbor attractive interaction, one obtains a similar Hamiltonian with $\Delta_y = -\frac{1}{3}(\Delta_{12} + \Delta_{13} + \Delta_{23})$, $\Delta_x = -\frac{1}{3}(2\Delta_{12} - \Delta_{13} - \Delta_{23})$, $\Delta_z = \frac{1}{3}(\Delta_{23} - \Delta_{13})$. Here $\Delta_{ij} = \frac{V'}{4} \sum_{\mathbf{p}} \langle c_{-\mathbf{p}i} c_{\mathbf{p}j} + c_{-\mathbf{p}j} c_{\mathbf{p}i} \rangle \sin p_{ij} \sin k_{ij}$ for $i, j = 1, 2, 3$, where V' is the strength of the attractive interaction. The pairing is then also *p*-wave type.

Projected BdG Hamiltonian and Chern number.— We focus on the weak pairing regime, where $|\mu|$ is much larger than $|\Delta_\nu(\mathbf{k})|$. In this regime one can safely ignore the coupling between states separated with energy difference $\geq |\mu|$. Consider $\mu > 0$ (the results are similar for $\mu < 0$ in the checkerboard lattice), where we can project the BdG Hamiltonian into the subspace with only the $+$ band which crosses the Fermi level. For simplicity, we consider the situation with $t_x = t_z$ in the checkerboard lattice. The spectrum is then $\varepsilon_{\mathbf{k}\pm} = k^2(t_0 \pm t_z)$ and $\varepsilon_{\mathbf{k}\pm} = \frac{1}{8}|t|k^2(1 \pm 1)$ in checkerboard and kagome lattices respectively. After the projection, one obtains the Hamiltonian $H_{\text{PBdG}} = \frac{1}{2} \sum_{\mathbf{k}} \Psi_P^\dagger(\mathbf{k}) \mathcal{H}_{\mathbf{k}}^P \Psi_P(\mathbf{k}) + \frac{1}{2} E_0$ with $\Psi_P(\mathbf{k}) = (c_{\mathbf{k}+}, c_{-\mathbf{k}+}^\dagger)^T$ and

$$\mathcal{H}_{\mathbf{k}}^P = \begin{bmatrix} \varepsilon_{\mathbf{k}+} - \mu & \Delta_{\text{eff}}(\mathbf{k}) \\ \Delta_{\text{eff}}^*(\mathbf{k}) & -\varepsilon_{\mathbf{k}+} + \mu \end{bmatrix}, \quad (6)$$

where

$$\Delta_{\text{eff}}(\mathbf{k}) = \Delta_y(\mathbf{k}) + \sum_{\pm} \frac{1}{2} e^{\pm 2i\theta_{\mathbf{k}}} [\Delta_x(\mathbf{k}) \mp i\Delta_z(\mathbf{k})]. \quad (7)$$

The spectrum is $\pm E_{\mathbf{k}}$ with $E_{\mathbf{k}} = \sqrt{\xi_{\mathbf{k}}^2 + |\Delta_{\text{eff}}(\mathbf{k})|^2}$ and $\xi_{\mathbf{k}} = \varepsilon_{\mathbf{k}+} - \mu$. One finds again that the singlet pairing does not contribute to $\Delta_{\text{eff}}(\mathbf{k})$ and the gap. The Chern number of the fully-gapped state is[13]

$$N_C = \frac{1}{2\pi} \int_0^{2\pi} d\theta_{\mathbf{k}} \partial_{\theta_{\mathbf{k}}} \theta_{\Delta}(\mathbf{k}) \Big|_{\text{FS}} \quad (8)$$

with $\theta_{\Delta}(\mathbf{k}) = \text{Arg}[\Delta_{\text{eff}}(\mathbf{k})]$, which is just the winding number of the effective pairing $\Delta_{\text{eff}}(\mathbf{k})$ at Fermi surface. A crucial observation is that the Chern number can only be *odd*, as only the triplet pairing contribute to $\Delta_{\text{eff}}(\mathbf{k})$. Hence *all* the fully-gapped pairing states are *non-Abelian* states[13].




FIG. 2. (Color online) In checkerboard lattice systems. (a) The ratio of the order parameters $|\Phi_1|/|\Phi_{-3}|$ as function of V/U . (b) Energy spectrum of the Bogoliubov quasi-particle versus k_y in a stripe with periodic (open) boundary condition along the y (x) -direction. The parameters are $t_0 = 10$, $t_z = t_x = 20$, $\mu = 2$, $V/2 = U = 1$, $g = 0.95$. The width of the stripe along x direction is $N_x = 501$ unit cells.

Non-Abelian ground states.— To determine the ground states we study the mean field free energy

$$F = -k_B T \sum_{\mathbf{k}} \ln \left(2 \cosh \frac{E_{\mathbf{k}}}{2k_B T} \right) + \frac{1}{2} E_0. \quad (9)$$

To facilitate the discussion, for checkerboard lattice we introduce $g_{\nu\beta} = \sqrt{\frac{V_\nu}{2}} \sum_{\mathbf{p}} p_{-\beta} \langle d_\nu^\dagger(\mathbf{p}) \rangle$ for $\beta = \pm$. This enables us to write $E_0 = g^2$ with $g = \sqrt{\sum_{\nu,\beta} |g_{\nu\beta}|^2}$ being the “pairing amplitude” and $\Delta_{\text{eff}}(\mathbf{k}) = \frac{k\sqrt{U}}{2\sqrt{2}} \sum_n \Phi_n e^{in\theta_{\mathbf{k}}}$ with $n = \pm 1, \pm 3$, where $\Phi_{\pm 1} = 2g_{y\pm} + g_{x\mp} \mp i\sqrt{\frac{V}{2U}} g_{z\mp}$, $\Phi_{\pm 3} = g_{x\pm} \mp i\sqrt{\frac{V}{2U}} g_{z\pm}$. By minimizing the free energy numerically, we find that there are two possible ground states with the same free energy: i) Φ_1 and Φ_{-3} are finite with $\Phi_{-1} = \Phi_3 = 0$, ii) Φ_{-1} and Φ_3 are finite with $\Phi_1 = \Phi_{-3} = 0$. Both two states are fully-gapped and break time-reversal symmetry. In fact, they are time-reversal partners. Calculation indicates that the Chern number of the two states are $N_C = \pm 1$ as $|\Phi_1| > |\Phi_{-3}|$ or $|\Phi_{-1}| > |\Phi_3|$. In Fig. 2(a), we plot the ratio $|\Phi_1|/|\Phi_{-3}|$ as a function of V/U . It is seen that $|\Phi_1| > |\Phi_{-3}|$ in all the parameter regime. At $V/U = 2$, $\Phi_{-3} = 0$ where $\Delta_y : \Delta_x : \Delta_z = \frac{2}{\sqrt{6}} : \frac{1}{\sqrt{6}} : i\frac{1}{\sqrt{6}}$. Within the mean field theory, for *all* V/U the ground state is a non-Abelian state with fully-gapped Fermi surface. We also plot the spectrum of the Bogoliubov quasi-particles in Fig. 2(b) at $V/U = 2$ as an illustration of the edge states. From the figure it is seen that there are two gapless chiral edge states which are localized at the two boundaries separately. The self-consistent equation for the pairing magnitude g at $V/U = 2$ is $1 = \frac{3}{8} \sum_{\mathbf{k}} \tanh(\frac{E_{\mathbf{k}}}{2k_B T}) E_{\mathbf{k}}^{-1} k^2 U$ with

$E_{\mathbf{k}} = \sqrt{\xi_{\mathbf{k}}^2 + \frac{3}{4}k^2 U g^2}$. From this we obtain the transition temperature $T_c \simeq \frac{2\gamma}{\pi} \sqrt{\Lambda \mu} \exp[-\frac{16\pi(t_0+t_z)^2}{3U\mu} + \frac{\Lambda-\mu}{2\mu}]$, where γ is the Euler constant and Λ is the high energy cut-off. In 2D the SC/SF phase transition is determined by the Kosterlitz-Thouless transition. However, in the weak coupling regime, the Kosterlitz-Thouless transition temperature is close to that obtained by the above approach[23, 24].

For kagome lattice, we find that the ground state is also a fully-gapped *non-Abelian* state where $\Delta_{12} : \Delta_{13} : \Delta_{23} = 1 : e^{i\pi/3} : e^{i2\pi/3}$ (only Φ_1 is finite and $N_C = 1$) or $\Delta_{12} : \Delta_{13} : \Delta_{23} = 1 : e^{-i\pi/3} : e^{-i2\pi/3}$ (only Φ_{-1} is nonzero, $N_C = -1$). The transition temperature is $T_c \simeq \frac{2\gamma}{\pi} \sqrt{\Lambda \mu} \exp[-\frac{3\pi|t|^2}{4V'\mu} + \frac{\Lambda-\mu}{2\mu}]$. In all those cases, the ground states are the fully-gapped states, which is partly due to that such states gain more condensation energy than the nodal states[15, 16].

Generalizations.— Here we generalize the property that *all* the singlet pairing instabilities are *absent* to more classes of multi-orbital systems. As Eq. (2) is a general description of two band touching, quite generally, the property comes from the fact that $\mathbf{h}(-\mathbf{k}) = \mathbf{h}(\mathbf{k})$. Hence the property can also be realized in time-reversal symmetric systems where the Fermi surface encloses a single two-band touching with any *even* winding number. Such two-band touching is stable in systems with time-reversal and space-inversion symmetry[13, 22]. Moreover, the property holds even when such band touching is gapped by finite $h_y(\mathbf{k})$ via time-reversal symmetry breaking, given that $h_y(-\mathbf{k}) = h_y(\mathbf{k})$ (space inversion symmetric). At finite $h_y(\mathbf{k})$ the property remains to be true even when $|h_0(\mathbf{k})| > \sqrt{h_x^2(\mathbf{k}) + h_z^2(\mathbf{k})}$ as long as there is only one band crosses the Fermi level and another is far away from it. It is also found[13] that in those systems *all* the fully-gapped states are *non-Abelian* states. In analog with the checkerboard and kagome lattice systems, the non-Abelian states are possibly the ground state in more general systems as they have fully-gapped Fermi surface to gain more condensation energy[15, 16].

Experimental realization and detections.— The checkerboard optical lattice has been realized in experiments[25]. The scheme to realize kagome optical lattice are proposed in Ref. [26]. The isotropic attractive interactions can be realized in Bose-Fermi mixtures in deep lattices[18] or in polar-molecule gases dressed by microwaves[19, 24]. Below we estimate the transition temperature for polar-molecule gases in checkerboard lattices. To have T_c in the experimentally observable range, a moderately strong interaction is needed[24]. We take $t_0 = 0$, $\Lambda = 6t_z$ (the band width is $15t_z$), (e.g.) $\mu = 0.8t_z$, and $U = 4t_z$, which gives $T_c \simeq 0.1t_z$ while the weak pairing condition $|\Delta_{\text{eff}}|_{T=0} \sim k_B T_c \ll |\mu|$ is still satisfied. For a lattice with $t_z = 200$ nK, $T_c \simeq 20$ nK. The required interaction (1600 nK=220 KHz) is in the experimentally accessi-

ble range for polar-molecules[19]. The pairing symmetry and the non-Abelian statistics in the proposed system can also be detected experimentally via recently developed/proposed techniques[7, 12, 20].

Acknowledgements.— I thank G. Conduit, S. Wu, J. Ruhman, M. Gong, and Y. E. Krauss for fruitful discussions and comments. This work was supported by the German Federal Ministry of Education and Research (BMBF) within the framework of the German-Israeli project cooperation (DIP) and by the Israel Science Foundation (ISF).

* jianhua.jiang@weizmann.ac.il

- [1] G. Moore and N. Read, Nucl. Phys. B **360**, 362 (1991).
- [2] A. Kitaev, Ann. Phys. **321**, 2 (2006).
- [3] N. Read and D. Green, Phys. Rev. B **61**, 10267 (2000).
- [4] D. A. Ivanov, Phys. Rev. Lett. **86**, 268 (2001).
- [5] M. Freedman *et al.*, Bull. Am. Math. Soc. **40**, 31 (2002); S. B. Bravyi and A. Y. Kitaev, Ann. Phys. (N.Y.) **298**, 210 (2002); A. Y. Kitaev, Ann. Phys. (N.Y.) **303**, 2 (2003).
- [6] C. Nayak *et al.*, Rev. Mod. Phys. **80**, 1083 (2008).
- [7] S. Tewari, S. Das Sarma, C. Nayak, C. Zhang, and P. Zoller, Phys. Rev. Lett. **98**, 010506 (2007).
- [8] X. G. Wen, *Quantum Field Theory of Many-Body Systems* (Oxford University Press, Oxford, 2004).
- [9] M. Sato, Phys. Lett. B **575**, 126 (2003); L. Fu and C. L. Kane, Phys. Rev. Lett. **100**, 096407 (2008).
- [10] C. Zhang, S. Tewari, R. M. Lutchyn, and S. Das Sarma, Phys. Rev. Lett. **101**, 160401 (2008); M. Sato, Y. Takahashi, and S. Fujimoto, *ibid.* **103**, 020401 (2009); M. Sato and S. Fujimoto, Phys. Rev. B **79**, 094504 (2009); J. D. Sau, *et al.*, Phys. Rev. Lett. **104**, 040502 (2010); J. Alicea, Phys. Rev. B **81**, 125318 (2010); R. M. Lutchyn, J. D. Sau, and S. Das Sarma, Phys. Rev. Lett. **105**, 077001 (2010); Y. Oreg, G. Refael, and F. von Oppen, *ibid.* **105**, 177002 (2010); L. Mao, J. Shi, Q. Niu, and C. Zhang, *ibid.* **106**, 157003 (2011).
- [11] X.-L. Qi, T. L. Hughes, and S. C. Zhang, Phys. Rev. B **82**, 184516 (2010).
- [12] S.-L. Zhu, L. B. Shao, Z. D. Wang, and L. M. Duan, Phys. Rev. Lett. **106**, 100404 (2011).
- [13] J. H. Jiang and S. Wu, arXiv:1107.4274
- [14] K. Sun, W. V. Liu, S. Das Sarma, arXiv:1011.4301
- [15] P. W. Anderson and P. Morel, Phys. Rev. **123**, 1911 (1961).
- [16] M. Cheng, K. Sun, V. Galitski, and S. Das Sarma, Phys. Rev. B **81**, 024504 (2010) [Note that this paper is focused on single band (orbital) systems].
- [17] C. Chin *et al.*, Rev. Mod. Phys. **82**, 1225 (2010).
- [18] M. Lewenstein, L. Santos, M. A. Baranov, and H. Fehrmann, Phys. Rev. Lett. **92**, 050401 (2004).
- [19] See, e.g., K.-K. Ni *et al.*, Science **322**, 231 (2008); J. Deiglmayr *et al.*, Phys. Rev. Lett. **101**, 133004 (2008); A. Micheli, G. K. Brennen, and P. Zoller, Nature Phys. **2**, 341 (2006); A. Micheli, G. Pupillo, H. P. Büchler, and P. Zoller, Phys. Rev. A **76**, 043604 (2007).
- [20] I. Bloch, J. Dalibard, and W. Zwerger, Rev. Mod. Phys. **80**, 885 (2008).

- [21] See, e.g., P. Soltan-Panahi *et al.*, Nature Phys. **7**, 434 (2011); *ibid.*, arXiv:1104.3456; W.-C. Lee, C. Wu, and S. Das Sarma, Phys. Rev. A **82**, 053611 (2010); H.-H. Hung, W.-C. Lee, C. Wu, Phys. Rev. B **83**, 144506 (2011).
- [22] K. Sun, H. Yao, E. Fradkin, and S. A. Kivelson, Phys. Rev. Lett. **103**, 046811 (2009).
- [23] K. Miyake, Prog. Theor. Phys. **69**, 1794 (1983).
- [24] N. R. Cooper and G. V. Shlyapnikov, Phys. Rev. Lett. **103**, 155302 (2009).
- [25] G. Wirth, M. Ölschläger, and A. Hemmerich, Nature Phys. **7**, 147 (2010); M. Ölschläger, G. Wirth, and A. Hemmerich, Phys. Rev. Lett. **106**, 015302 (2011).
- [26] J. Ruostekoski, Phys. Rev. Lett. **103**, 080406 (2009).